

**Heterogeneous Variance Models from Hoffman Chapter 8:
Predicting Differential Variation in Fluctuation of Daily Physical Symptoms from
Sex, Age, and Daily Negative Mood (PMC) and Stressor Days (GMC)**

Same baseline homogeneous variance model via SAS PROC MIXED or via SAS PROC NLMIXED:

```

TITLE1 "Baseline Homogeneous Variance Model with just Fixed Effects";
PROC MIXED DATA=example COVTEST NOCLPRINT NOITPRINT NOIINFO NAMELEN=50 METHOD=ML;
CLASS ID;
MODEL symptom5 = female age80 female*age80 PMmood2 PMstress40 female*PMstress40 PMmood2*PMmood2
               WPmood PMmood2*WPmood TVstress0 female*TVstress0 / SOLUTION DDFM=SATTERTHWAITE;
RANDOM INTERCEPT / TYPE=UN SUBJECT=ID;
REPEATED / TYPE=VC SUBJECT=ID; RUN; TITLE1;

TITLE1 "Baseline Homogeneous Variance Model with just Fixed Effects";
PROC NLMIXED DATA=example METHOD=GAUSS TECH=NEWRAP GCONV=1e-12;
* Must define all parameters to be estimated and provide start values;
* First line is fixed effects, second line is variances;
PARMS fint=.245 fsex=-.33 fage=.07 fsexage=-.09
      fbpmood=-.34 fwpmood=.26 fcstress=1.93 fwptress=.22
      f2sexstress=-1.33 f1sexstress=-.18 fbpmood2=-1.78 fbpwpwmood=.26
      vu0int=-.47 varEint=-.50;
* Setting up level-2 equations;
b0i = fint + fsex*female + fage*age80 + fsexage*female*age80
      + fbpmood*PMmood2 + fcstress*PMstress40
      + f2sexstress*PMstress40*female + fbpmood2*PMmood2*PMmood2 + u0i;
b1i = fwpmood + fbpwpwmood*PMmood2;
b2i = fwptress + f1sexstress*female;
* Setting up level-1 equation WITHOUT level-1 residual;
y = (b0i) + (b1i*WPmood) + (b2i*TVstress0);
* Model for heterogeneous variance at level-2 and level-1;
vu0 = EXP(vu0int);
varE = EXP(varEint);
* Telling it which DV, defining level-1 residual;
* symptom5 is normally distributed with a mean of "y" and a variance of "VarE";
      MODEL symptom5 ~ normal(y,varE);
* Defining random effects: normally distributed with means and variances;
      RANDOM u0i ~ normal([0],[vu0]) SUBJECT=ID;
* Requesting additional model-implied estimates;
      ESTIMATE "Actual Random Intercept Variance" EXP(vu0int);
      ESTIMATE "Actual Residual Variance"           EXP(varEint);
RUN; TITLE1;

```

Level 1:

$$\text{Symptoms}_{ti} = \beta_{0i} + \beta_{1i} (\text{Mood}_{ti} - \overline{\text{Mood}}_i) + \beta_{2i} (\text{Stressor}_{ti}) + e_{ti}$$

$$\text{Residual Variance: } \sigma_{e_{ti}}^2 = \exp[\eta_{0i}]$$

Level 2 :

$$\begin{aligned} \text{Intercept: } \beta_{0i} &= \gamma_{00} + \gamma_{01} (\text{Women}_i) + \gamma_{02} (\text{Age}_i - 80) + \gamma_{03} (\text{Women}_i)(\text{Age}_i - 80) + \gamma_{04} (\text{Mood}_i - 2) \\ &\quad + \gamma_{08} (\overline{\text{Stressor}}_i - 0.40) + \gamma_{09} (\text{Women}_i)(\overline{\text{Stressor}}_i - 0.40) + \gamma_{0,16} (\overline{\text{Mood}}_i - 2)^2 + U_{0i} \end{aligned}$$

$$\text{Within-Person Mood: } \beta_{1i} = \gamma_{10} + \gamma_{14} (\overline{\text{Mood}}_i - 2)$$

$$\text{Within-Person Stressor: } \beta_{2i} = \gamma_{20} + \gamma_{21} (\text{Women}_i)$$

$$\text{Random Intercept Variance } \tau_{U_{0i}}^2 = \exp[\nu_{00}]$$

$$\text{Residual Variance: } \eta_{0i} = \varepsilon_{00}$$

PROC MIXED OUTPUT:

Covariance Parameter Estimates						
Cov Parm	Subject	Estimate	Standard Error	Value	Z	Pr > Z
UN(1,1)	id	0.6230	0.1044	5.96	<.0001	Random Intercept Variance
Residual	id	0.6093	0.04292	14.19	<.0001	Residual Variance
Information Criteria						
Neg2LogLike	Parms	AIC	AICC	HQIC	BIC	CAIC
1379.2	14	1407.2	1408.1	1422.3	1444.4	1458.4
Solution for Fixed Effects						
Effect	Estimate	Error	DF	t Value	Pr > t	Standard
Intercept	2.4473	0.3474	117	7.05	<.0001	
female	-0.3277	0.2140	146	-1.53	0.1278	
age80	0.07311	0.03019	109	2.42	0.0171	
female*age80	-0.09127	0.03381	108	-2.70	0.0081	
PMmood2	-0.3396	0.7999	110	-0.42	0.6720	
PMstress40	1.9322	0.6005	124	3.22	0.0016	
female*PMstress40	-1.3281	0.6617	128	-2.01	0.0468	
PMmood2*PMmood2	-1.7784	0.7131	108	-2.49	0.0141	
WPmood	0.2634	0.2213	405	1.19	0.2346	
PMmood2*WPmood	0.2591	0.3962	405	0.65	0.5135	
TVstress0	0.2192	0.1710	404	1.28	0.2007	
female*TVstress0	-0.1812	0.2053	404	-0.88	0.3780	

PROC NL MIXED OUTPUT:

Dimensions	
Observations Used	509
Observations Not Used	0
Total Observations	509
Subjects	105
Max Obs Per Subject	5
Parameters	14
Quadrature Points	1

Specifications	
Data Set	WORK.EXAMPLE
Dependent Variable	symptom5
Distribution for Dependent Variable	Normal
Random Effects	U0i
Distribution for Random Effects	Normal
Subject Variable	id
Optimization Technique	Newton-Raphson
Integration Method	Adaptive Gaussian Quadrature

NOTE: ABSGCONV convergence criterion satisfied.

Fit Statistics	
-2 Log Likelihood	1379.2
AIC (smaller is better)	1407.2
AICC (smaller is better)	1408.1
BIC (smaller is better)	1444.4

Make sure the gradient (partial first derivative with respect to each parameter) is as close to 0 as possible. Otherwise, the estimation really hasn't finished on the "best" estimates.

Parameter	Estimate	Error	DF	t Value	Pr > t	Parameter Estimates			
						Standard	Alpha	Lower	Upper
fint	2.4473	0.3474	104	7.05	<.0001	0.05	1.7585	3.1362	-6.09E-8
fsex	-0.3277	0.2140	104	-1.53	0.1286	0.05	-0.7520	0.09659	3.344E-8
fage	0.07311	0.03019	104	2.42	0.0172	0.05	0.01324	0.1330	5.738E-6
fsexage	-0.09127	0.03381	104	-2.70	0.0081	0.05	-0.1583	-0.02423	3.875E-6
fbpmood	-0.3396	0.7999	104	-0.42	0.6720	0.05	-1.9257	1.2466	1.156E-7
fpwmood	0.2634	0.2213	104	1.19	0.2366	0.05	-0.1754	0.7022	-1.09E-8
fcstress	1.9322	0.6005	104	3.22	0.0017	0.05	0.7415	3.1230	1.034E-8
fpwstress	0.2192	0.1710	104	1.28	0.2028	0.05	-0.1199	0.5584	-6.11E-9
f2sexstress	-1.3281	0.6617	104	-2.01	0.0473	0.05	-2.6402	-0.01600	1.161E-8
f1sexstress	-0.1812	0.2053	104	-0.88	0.3796	0.05	-0.5882	0.2259	3.322E-8
fbpmood2	-1.7784	0.7131	104	-2.49	0.0142	0.05	-3.1926	-0.3643	-9.03E-8
fbpwpwmood	0.2591	0.3962	104	0.65	0.5145	0.05	-0.5266	1.0449	7.878E-9
vU0int	-0.4733	0.1677	104	-2.82	0.0057	0.05	-0.8057	-0.1408	-5.58E-7
varEint	-0.4955	0.07045	104	-7.03	<.0001	0.05	-0.6352	-0.3558	-7.07E-7

Label	Estimate	Additional Estimates						
		Standard	Error	DF	t Value	Pr > t	Alpha	Lower
Actual Random Intercept Variance	0.6230	0.1044	104	5.96	<.0001	0.05	0.4159	0.8301
Actual Residual Variance	0.6093	0.04292	104	14.19	<.0001	0.05	0.5241	0.6944

Testing a ω_{0i} scale factor for random individual differences in within-person variation:

Level 1 :

$$\text{Symptoms}_{ti} = \beta_{0i} + \beta_{1i} (\text{Mood}_{ti} - \overline{\text{Mood}}_i) + \beta_{2i} (\text{Stressor}_{ti}) + \epsilon_{ti}$$

$$\text{Residual Variance: } \sigma_{\epsilon_{ti}}^2 = \exp[\eta_{0i}]$$

Level 2 :

$$\begin{aligned} \text{Intercept: } \beta_{0i} &= \gamma_{00} + \gamma_{01} (\text{Women}_i) + \gamma_{02} (\text{Age}_i - 80) + \gamma_{03} (\text{Women}_i)(\text{Age}_i - 80) + \gamma_{04} (\overline{\text{Mood}}_i - 2) \\ &\quad + \gamma_{08} (\overline{\text{Stressor}}_i - 0.40) + \gamma_{09} (\text{Women}_i)(\overline{\text{Stressor}}_i - 0.40) + \gamma_{0,16} (\overline{\text{Mood}}_i - 2)^2 + U_{0i} \end{aligned}$$

$$\text{Within-Person Mood: } \beta_{1i} = \gamma_{10} + \gamma_{14} (\overline{\text{Mood}}_i - 2)$$

$$\text{Within-Person Stressor: } \beta_{2i} = \gamma_{20} + \gamma_{21} (\text{Women}_i)$$

$$\text{Random Intercept Variance } \tau_{U_{0i}}^2 = \exp[\nu_{00}]$$

$$\text{Residual Variance: } \eta_{0i} = \epsilon_{00} + \omega_{0i}$$

```

TITLE1 "Baseline Model with just Fixed Effects";
TITLE2 "Test Scale Factor";
PROC NLINMIXED DATA=example METHOD=GAUSS TECH=NEWRAP GCONV=1e-12; *METHOD=FIRO;
* Must define all parameters to be estimated and provide start values;
* First line is fixed effects, second line is variances;
PARMS fint=2.45 fsex=-.33 fage=.07 fsexage=-.09
      fbpmmood=-.34 fwpmood=.26 fcstress=1.93 fwpmstress=.22
      f2sexstress=-1.33 f1sexstress=-.18 fbpmmood2=-1.78 fbpwpmood=.26
      varEint=-.50 vU0int=-.47 vscale=-.22 cU0scale=.00;
* Setting up level-2 equations;
b0i = fint + fsex*female + fage*age80 + fsexage*female*age80
      + fbpmmood*PMmood2 + fcstress*PMstress40
      + f2sexstress*PMstress40*female + fbpmmood2*PMmood2*PMmood2 + U0i;
b1i = fwpmood + fbpwpmood*PMmood2;
b2i = fwpmstress + f1sexstress*female;
* Setting up level-1 equation WITHOUT level-1 residual;
y = (b0i) + (b1i*WPmood) + (b2i*TVstress0);
* Model for heterogeneous variance at level-2 and level-1;
vU0 = EXP(vU0int);
varE = EXP(varEint + Scalei);
* Telling it which DV, defining level-1 residual;
* symptom5 is normally distributed with a mean of "y" and a variance of "VarE";
      MODEL symptom5 ~ normal(y, varE);
* Defining random effects: normally distributed with means and variances;
      RANDOM U0i Scalei ~ normal([0,0],[vU0,cU0scale,vscale]) SUBJECT=ID;
RUN; TITLE1; TITLE2;

```

PROC NLINMIXED OUTPUT:

Specifications	
Data Set	WORK.EXAMPLE
Dependent Variable	symptom5
Distribution for Dependent Variable	Normal
Random Effects	U0i Scalei
Distribution for Random Effects	Normal
Subject Variable	id
Optimization Technique	Newton-Raphson
Integration Method	Adaptive Gaussian Quadrature

That's all we get – SAS could not find an estimate for the scale factor variance, even after switching to less intensive estimation routines (e.g., METHOD=FIRO), so we must proceed assuming we don't have one.

Examining predictors of differential level-2 random intercept variance across persons:

Level 1 :

$$\text{Symptoms}_{ti} = \beta_{0i} + \beta_{1i} (\text{Mood}_{ti} - \overline{\text{Mood}}_i) + \beta_{2i} (\text{Stressor}_{ti}) + e_{ti}$$

$$\text{Residual Variance: } \sigma_{e_{ti}}^2 = \exp[\eta_{0i}]$$

Level 2 :

$$\begin{aligned} \text{Intercept: } \beta_{0i} &= \gamma_{00} + \gamma_{01} (\text{Women}_i) + \gamma_{02} (\text{Age}_i - 80) + \gamma_{03} (\text{Women}_i)(\text{Age}_i - 80) + \gamma_{04} (\overline{\text{Mood}}_i - 2) \\ &\quad + \gamma_{08} (\overline{\text{Stressor}}_i - 0.40) + \gamma_{09} (\text{Women}_i)(\overline{\text{Stressor}}_i - 0.40) + \gamma_{0,16} (\overline{\text{Mood}}_i - 2)^2 + U_{0i} \end{aligned}$$

$$\text{Within-Person Mood: } \beta_{1i} = \gamma_{10} + \gamma_{14} (\overline{\text{Mood}}_i - 2)$$

$$\text{Within-Person Stressor: } \beta_{2i} = \gamma_{20} + \gamma_{21} (\text{Women}_i)$$

$$\text{Random Intercept Variance } \tau_{U_{0i}}^2 = \exp \left[\begin{array}{l} v_{00} + v_{01}(\text{Women}_i) + v_{02}(\text{Age}_i - 80) \\ + v_{01}(\overline{\text{Mood}}_i - 2) + v_{08}(\overline{\text{Stressor}}_i - 0.40) \end{array} \right]$$

$$\text{Residual Variance: } \eta_{0i} = \varepsilon_{00}$$

```

TITLE1 "Add Level 2 Predictors for Sex, Age, Mood, and Stress";
PROC NLIN MIXED DATA=example METHOD=GAUSS TECH=NEWRAP GCONV=1e-12;
* Must define all parameters to be estimated and provide start values;
* First line is fixed effects, second line is variances;
PARMS fint=2.45 fsex=-.33 fage=.07 fsexage=-.09
      fbpmood=-.34 fwpmood=.26 fcstress=1.93 fwpmstress=.22
      f2sexstress=-1.33 f1sexstress=-.18 fbpmood2=-1.78 fbwpwmood=.26
      vu0int=-.47 vu0sex=0 vu0age=0 vu0bpwmood=0 vu0bpstress=0 varEint=-.50;
* Setting up level-2 equations;
b0i = fint + fsex*female + fage*age80 + fsexage*female*age80
      + fbpmood*PMmood2 + fcstress*PMstress40
      + f2sexstress*PMstress40*female + fbpmood2*PMmood2*PMmood2 + u0i;
b1i = fwpmood + fbwpwmood*PMmood2;
b2i = fwpmstress + f1sexstress*female;
* Setting up level-1 equation WITHOUT level-1 residual;
y = (b0i) + (b1i*WPmood) + (b2i*TVstress0);
* Model for heterogeneous variance at level-2 and level-1;
vu0 = EXP(vu0int + vu0sex*female + vu0age*age80 + vu0bpwmood*PMmood2 + vu0bpstress*PMstress40);
varE = EXP(varEint);
* Telling it which DV, defining level-1 residual;
* symptom5 is normally distributed with a mean of "y" and a variance of "VarE";
      MODEL symptom5 ~ normal(y,varE);
* Defining random effects: normally distributed with means and variances;
      RANDOM u0i ~ normal([0],[vu0]) SUBJECT=id;
RUN; TITLE1;

```

		Specifications
Dimensions		
Observations Used	509	WORK.EXAMPLE
Observations Not Used	0	symptom5
Total Observations	509	Normal
Subjects	105	U0i
Max Obs Per Subject	5	Normal
Parameters	18	id
Quadrature Points	1	Newton-Raphson
		Adaptive Gaussian Quadrature

NOTE: GCONV convergence criterion satisfied.

Fit Statistics	
-2 Log Likelihood	1371.0
AIC (smaller is better)	1407.0
AICC (smaller is better)	1408.4
BIC (smaller is better)	1454.8

The previous model was $-2\text{LL}(14) = 1379.2$.
The current model is $-2\text{LL}(18) = 1371.0$, so
 $-2\Delta\text{LL}(4) = 8.2$, $p = .085$, although only one new effect is significant individually...

Parameter	Parameter Estimates								
	Estimate	Error	DF	t Value	Pr > t	Alpha	Lower	Upper	Gradient
fint	2.4208	0.3426	104	7.07	<.0001	0.05	1.7415	3.1001	-474E-12
fsex	-0.3364	0.2099	104	-1.60	0.1121	0.05	-0.7527	0.07992	-505E-12
fage	0.07753	0.03087	104	2.51	0.0136	0.05	0.01631	0.1387	4.143E-9
fsexage	-0.08882	0.03427	104	-2.59	0.0109	0.05	-0.1568	-0.02086	3.719E-9
fbpmood	-0.2178	0.7821	104	-0.28	0.7812	0.05	-1.7688	1.3332	4.97E-10
fpwmood	0.2616	0.2212	104	1.18	0.2396	0.05	-0.1770	0.7002	2.09E-12
fcstress	1.8901	0.6084	104	3.11	0.0024	0.05	0.6837	3.0965	1.18E-10
fpstress	0.2233	0.1710	104	1.31	0.1945	0.05	-0.1157	0.5623	-112E-12
f2sexstress	-1.2330	0.6696	104	-1.84	0.0684	0.05	-2.5608	0.09487	1.08E-10
f1sexstress	-0.1850	0.2052	104	-0.90	0.3692	0.05	-0.5919	0.2218	-107E-12
fbpmood2	-1.5905	0.6760	104	-2.35	0.0205	0.05	-2.9310	-0.2501	-372E-12
fbpwpmmood	0.2564	0.3960	104	0.65	0.5188	0.05	-0.5289	1.0416	1.77E-13
vU0int	-1.1082	0.7895	104	-1.40	0.1634	0.05	-2.6739	0.4575	-215E-12
vU0sex	-0.1760	0.3852	104	-0.46	0.6487	0.05	-0.9398	0.5878	-211E-12
vU0age	0.03867	0.03293	104	1.17	0.2430	0.05	-0.02663	0.1040	6.8E-10
vU0bpmmood	-0.7150	0.8469	104	-0.84	0.4004	0.05	-2.3944	0.9644	1.71E-10
vU0bpstress	1.5833	0.6031	104	2.63	0.0100	0.05	0.3874	2.7792	2.87E-11
varEint	-0.4968	0.07034	104	-7.06	<.0001	0.05	-0.6363	-0.3573	-133E-12

The effect of PMstressor40_i on the amount of random intercept variance indicates that for every unit higher proportion of stressor days, the log of the random intercept variance is predicted to be higher by 1.58. In other words, there is significantly greater person-to-person variation in mean daily reported physical symptoms among people who report experiencing a greater proportion of stressor days.

Examining predictors of differential level-1 residual variance across persons (retaining all effects of heterogeneity on the level-2 random intercept variance for now):

Level 1:

$$\text{Symptoms}_{ti} = \beta_{0i} + \beta_{1i} (\text{Mood}_{ti} - \overline{\text{Mood}}_i) + \beta_{2i} (\text{Stressor}_{ti}) + e_{ti}$$

$$\text{Residual Variance: } \sigma_{e_{ti}}^2 = \exp \left[\eta_{0i} + \eta_{1i} (\text{Mood}_{ti} - \overline{\text{Mood}}_i) + \eta_{2i} (\text{Stressor}_{ti}) \right]$$

Level 2:

$$\begin{aligned} \text{Intercept: } \beta_{0i} &= \gamma_{00} + \gamma_{01} (\text{Women}_i) + \gamma_{02} (\text{Age}_i - 80) + \gamma_{03} (\text{Women}_i)(\text{Age}_i - 80) + \gamma_{04} (\overline{\text{Mood}}_i - 2) \\ &\quad + \gamma_{08} (\overline{\text{Stressor}}_i - 0.40) + \gamma_{09} (\text{Women}_i)(\overline{\text{Stressor}}_i - 0.40) + \gamma_{0,16} (\overline{\text{Mood}}_i - 2)^2 + U_{0i} \end{aligned}$$

$$\text{Within-Person Mood: } \beta_{1i} = \gamma_{10} + \gamma_{14} (\overline{\text{Mood}}_i - 2)$$

$$\text{Within-Person Stressor: } \beta_{2i} = \gamma_{20} + \gamma_{21} (\text{Women}_i)$$

$$\text{Random Intercept Variance } \tau_{U_{0i}}^2 = \exp \left[\begin{aligned} &\nu_{00} + \nu_{01} (\text{Women}_i) + \nu_{02} (\text{Age}_i - 80) \\ &+ \nu_{04} (\overline{\text{Mood}}_i - 2) + \nu_{08} (\overline{\text{Stressor}}_i - 0.40) \end{aligned} \right]$$

Residual Variance:

$$\eta_{0i} = \varepsilon_{00} + \varepsilon_{01} (\text{Women}_i) + \varepsilon_{02} (\text{Age}_i - 80) + \varepsilon_{04} (\overline{\text{Mood}}_i - 2) + \varepsilon_{08} (\overline{\text{Stressor}}_i - 0.40)$$

$$\eta_{1i} = \varepsilon_{10}$$

$$\eta_{2i} = \varepsilon_{20}$$

```

TITLE1 "Level-1 Heterogeneity by L2 for Sex, Age, Mood, and Stress, L1 Mood and Stress";
PROC NLIN MIXED DATA=example METHOD=GAUSS TECH=NEWRAP GCONV=1e-12;
* Must define all parameters to be estimated and provide start values;
* First line is fixed effects, second line is variances;
PARMS fint=.245 fsex=-.33 fage=.07 fsexage=-.09
      fbp mood=-.34 fwp mood=.26 fc stress=1.93 fwp stres=.22
      f2 sex stress=-1.33 f1 sex stress=-.18 fbp mood2=-1.78 fbp wp mood=.26
      vU0 int=-.47 vU0 sex=0 vU0 age=0 vU0 bp mood=0 vU0 bp stres=0
      var Eint=-.50 var Esex=0 var Eage=0 var Ebpmood=0 var Ebps tress=0 var Ewp mood=0 var Ewp stres=0;
* Setting up level-2 equations;
b0i = fint + fsex*female + fage*age80 + fsexage*female*age80
      + fbp mood*PMmood2 + fc stress*PMstress40
      + f2 sex stress*PMstress40*female + fbp mood2*PMmood2*PMmood2 + U0i;
b1i = fwp mood + fbp wp mood*PMmood2;
b2i = fwp stres + f1 sex stress*female;
* Setting up level-1 equation WITHOUT level-1 residual;
y = (b0i) + (b1i*WPmood) + (b2i*TVstress0);
* Model for heterogeneous variance at level-2 and level-1;
vU0 = EXP(vU0 int + vU0 sex*female + vU0 age*age80 + vU0 bp mood*PMmood2 + vU0 bp stres*PMstress40);
var E = EXP(var Eint + var Esex*female + var Eage*age80 + var Ebpmood*PMmood2
      + var Ebps tress*PMstress40 + var Ewp mood*WPmood + var Ewp stres*TVstress0);
* Telling it which DV, defining level-1 residual;
* symptom5 is normally distributed with a mean of "y" and a variance of "VarE";
      MODEL symptom5 ~ normal(y, var E);
* Defining random effects: normally distributed with means and variances;
      RANDOM U0i ~ normal([0], [vU0]) SUBJECT=ID;
RUN; TITLE1;

```

Dimensions

Observations Used	509
Observations Not Used	0
Total Observations	509
Subjects	105
Max Obs Per Subject	5
Parameters	24
Quadrature Points	1

NOTE: ABSGCONV convergence criterion satisfied.

Fit Statistics	
-2 Log Likelihood	1346.2
AIC (smaller is better)	1394.2
AICC (smaller is better)	1396.7
BIC (smaller is better)	1457.9

The previous model was $-2\text{LL}(18) = 1371.0$.
The current model is $-2\text{LL}(24) = 1346.2$, so
 $-2\Delta\text{LL}(6) = 24.8, p < .001$, although only two
new effects are significant individually...

Parameter	Estimate	Standard	Parameter Estimates						
			DF	t Value	Pr > t	Alpha	Lower	Upper	Gradient
fint	2.4319	0.3651	104	6.66	<.0001	0.05	1.7079	3.1560	-1.49E-7
fsex	-0.3358	0.2079	104	-1.62	0.1093	0.05	-0.7481	0.07647	-1.43E-7
fage	0.07538	0.03087	104	2.44	0.0163	0.05	0.01417	0.1366	8.042E-7
fsexage	-0.08680	0.03438	104	-2.52	0.0131	0.05	-0.1550	-0.01862	7.225E-7
fbpmood	-0.1622	0.8895	104	-0.18	0.8556	0.05	-1.9262	1.6017	1.228E-7
fpwmood	0.1526	0.3170	104	0.48	0.6311	0.05	-0.4759	0.7812	2.418E-8
fc stress	1.9231	0.5954	104	3.23	0.0017	0.05	0.7425	3.1037	2.812E-8
fwp stress	0.2081	0.1610	104	1.29	0.1989	0.05	-0.1111	0.5273	-2.94E-8
f2 sex stress	-1.2523	0.6588	104	-1.90	0.0601	0.05	-2.5586	0.05405	2.74E-8
f1 sex stress	-0.1748	0.1950	104	-0.90	0.3721	0.05	-0.5616	0.2120	-2.72E-8
fbpmood2	-1.5408	0.7463	104	-2.06	0.0415	0.05	-3.0208	-0.06075	-1.08E-7
fbp wp mood	0.08266	0.5216	104	0.16	0.8744	0.05	-0.9517	1.1170	-2E-9
vU0 int	-1.0869	0.8058	104	-1.35	0.1803	0.05	-2.6848	0.5110	-3.49E-8
vU0 sex	-0.1516	0.3839	104	-0.39	0.6938	0.05	-0.9129	0.6098	-3.36E-8
vU0 age	0.03924	0.03203	104	1.22	0.2234	0.05	-0.02428	0.1028	2.059E-7
vU0 bp mood	-0.7041	0.8717	104	-0.81	0.4211	0.05	-2.4327	1.0245	2.874E-8
vU0 bp stres	1.4262	0.5871	104	2.43	0.0169	0.05	0.2619	2.5904	6.579E-9
var Eint	-0.08445	0.3079	104	-0.27	0.7844	0.05	-0.6949	0.5260	-6.25E-8
var Esex	0.07127	0.1663	104	0.43	0.6691	0.05	-0.2585	0.4010	-4.75E-8
var Eage	-0.00337	0.01311	104	-0.26	0.7979	0.05	-0.02937	0.02264	-4.53E-8
var Ebpmood	0.6972	0.3368	104	2.07	0.0409	0.05	0.02922	1.3651	2.565E-8
var Ebps tress	0.6967	0.3152	104	2.21	0.0293	0.05	0.07160	1.3219	-4.66E-9
var Ewp mood	0.02650	0.3095	104	0.09	0.9319	0.05	-0.5873	0.6403	6.281E-9
var Ewp stres	-0.00774	0.2177	104	-0.04	0.9717	0.05	-0.4395	0.4241	-2.94E-8

The effect of PMmood2_i on the amount of residual variance indicates that for every unit higher person mean negative mood, the log of the residual variance is predicted to be higher by 0.6972. Likewise, the effect of PMstressor40_i on the amount of residual variance indicates that for every unit higher proportion of stressor days, the log of the residual variance is predicted to be higher by 0.6967. In other words, there is significantly greater within-person daily fluctuation in reported physical symptoms among people who report greater average levels of negative and who report experiencing a greater proportion of stressor days.

Best model (after trimming any effects that were no longer significant, as shown in Table 8.7):

Level 1:

$$\text{Symptoms}_{ti} = \beta_{0i} + \beta_{1i} (\text{Mood}_{ti} - \overline{\text{Mood}}_i) + \beta_{2i} (\text{Stressor}_{ti}) + e_{ti}$$

$$\text{Residual Variance: } \sigma_{e_{ti}}^2 = \exp[\eta_{0i}]$$

Level 2:

$$\begin{aligned} \text{Intercept: } \beta_{0i} &= \gamma_{00} + \gamma_{01} (\text{Women}_i) + \gamma_{02} (\text{Age}_i - 80) + \gamma_{03} (\text{Women}_i)(\text{Age}_i - 80) + \gamma_{04} (\overline{\text{Mood}}_i - 2) \\ &\quad + \gamma_{08} (\overline{\text{Stressor}}_i - 0.40) + \gamma_{09} (\text{Women}_i)(\overline{\text{Stressor}}_i - 0.40) + U_{0i} \end{aligned}$$

$$\text{Within-Person Mood: } \beta_{1i} = \gamma_{10}$$

$$\text{Within-Person Stressor: } \beta_{2i} = \gamma_{20} + \gamma_{21} (\text{Women}_i)$$

$$\text{Random Intercept Variance } \tau_{U_{0i}}^2 = \exp[v_{00} + v_{08} (\overline{\text{Stressor}}_i - 0.40)]$$

Residual Variance:

$$\eta_{0i} = \varepsilon_{00} + \varepsilon_{04} (\overline{\text{Mood}}_i - 2) + \varepsilon_{08} (\overline{\text{Stressor}}_i - 0.40)$$

```
TITLE1 "Final Heterogeneous Variance Model without Quad PM Mood and BP*WP Mood";
PROC NLIN MIXED DATA=example METHOD=GAUSS TECH=NEWRAP GCONV=1e-12;
* Must define all parameters to be estimated and provide start values;
* First line is fixed effects, second line is variances;
PARMS fint=2.45 fsex=-.33 fage=.07 fsexage=-.09
      fbpmood=1.66 fwpmood=.26
      fcstress=1.93 fwpstress=.22 f2sexstress=-1.33 f1sexstress=-.18
      vU0int=-.47 vU0bpstress=0
      varEint=-.50 varEbpmood=0 varEbpstress=0;

* Setting up level-2 equations;
b0ifixed = fint + fsex*female + fage*age80 + fsexage*female*age80
            + fbpmood*PMmood2 + fcstress*PMstress40
            + f2sexstress*PMstress40*female;
b1i = fwpmood;
b2i = fwpstress + f1sexstress*female;

* Setting up level-1 equation WITHOUT level-1 residual;
yfixed = (b0ifixed) + (b1i*WPmood) + (b2i*TVstress0);
y = yfixed + U0i;

* Model for heterogeneous variance at level-2 and level-1;
vU0 = EXP(vU0int + vU0bpstress*PMstress40);
varE = EXP(varEint + varEbpmood*PMmood2 + varEbpstress*PMstress40);

* Telling it which DV, defining level-1 residual;
* symptom5 is normally distributed with a mean of "y" and a variance of "VarE";
      MODEL symptom5 ~ normal(y, varE);
* Defining random effects: normally distributed with means and variances;
      RANDOM U0i ~ normal([0], [vU0]) SUBJECT=ID;
```

```

* Asking for fixed effect predicted values;
  PREDICT yfixed OUT=Predicted;
* Getting all model implied estimates;
  ESTIMATE "Intercept: Men"          fint;
  ESTIMATE "Intercept: Women"        fint + fsex;
  ESTIMATE "Intercept: Sex Diff"     fsex;
  ESTIMATE "Age Slope: Men"         fage;
  ESTIMATE "Age Slope: Women"       fage + fsexage;
  ESTIMATE "Age Slope: Sex Diff"    fsexage;
  ESTIMATE "BP Mood: Linear"       fbp mood;
  ESTIMATE "Contextual Mood: Linear" fbp mood - fwpm mood;
  ESTIMATE "WP Mood"               fwpm mood;
  ESTIMATE "BP Stress: Men"        fc stress + fwps stress;
  ESTIMATE "BP Stress: Women"      fc stress + f2sex stress + fwps stress + f1sex stress;
  ESTIMATE "BP Stress: Sex Diff"   f2sex stress + f1sex stress;
  ESTIMATE "Contextual Stress: Men" fc stress;
  ESTIMATE "Contextual Stress: Women" fc stress + f2sex stress;
  ESTIMATE "Contextual Stress: Sex Diff" f2sex stress;
  ESTIMATE "WP Stress: Men"        fwps stress;
  ESTIMATE "WP Stress: Women"      fwps stress + f1sex stress;
  ESTIMATE "WP Stress: Sex Diff"   f1sex stress;
  ESTIMATE "Contextual Stress per Day: Men" fc stress/5;
  ESTIMATE "Contextual Stress per Day: Women" (fc stress + f2sex stress)/5;
RUN; PROC CORR DATA=Predicted; VAR symptom5 pred; RUN; TITLE1;

```

Specifications

Data Set	WORK.EXAMPLE
Dependent Variable	symptom5
Distribution for Dependent Variable	Normal
Random Effects	U0i
Distribution for Random Effects	Normal
Subject Variable	id
Optimization Technique	Newton-Raphson
Integration Method	Adaptive Gaussian Quadrature

Dimensions

Observations Used	509
Observations Not Used	0
Total Observations	509
Subjects	105
Max Obs Per Subject	5
Parameters	15
Quadrature Points	1

NOTE: ABSGCONV convergence criterion satisfied.

Fit Statistics

-2 Log Likelihood	1351.8
AIC (smaller is better)	1381.8
AICC (smaller is better)	1382.7
BIC (smaller is better)	1421.6

Parameter Estimates

Parameter	Estimate	Error	DF	t Value	Pr > t	Alpha	Lower	Upper	Gradient
fint	2.8017	0.3680	104	7.61	<.0001	0.05	2.0719	3.5314	4.992E-6
fsex	-0.3413	0.2059	104	-1.66	0.1004	0.05	-0.7495	0.06699	3.998E-6
fage	0.06876	0.03007	104	2.29	0.0243	0.05	0.009126	0.1284	-1.21E-6
fsexage	-0.08837	0.03327	104	-2.66	0.0092	0.05	-0.1543	-0.02239	4.658E-7
fbpmood	1.6584	0.3974	104	4.17	<.0001	0.05	0.8703	2.4464	-4.42E-6
fwpmood	0.1024	0.1473	104	0.69	0.4887	0.05	-0.1898	0.3945	-2.69E-9
fcstress	2.0496	0.6011	104	3.41	0.0009	0.05	0.8576	3.2415	-1.48E-6
fwpstress	0.2043	0.1647	104	1.24	0.2176	0.05	-0.1223	0.5309	5.092E-7
f2sexstress	-1.3699	0.6707	104	-2.04	0.0436	0.05	-2.6998	-0.03991	-1.24E-6
f1sexstress	-0.1754	0.1980	104	-0.89	0.3776	0.05	-0.5680	0.2171	3.609E-7
vU0int	-0.5724	0.1712	104	-3.34	0.0011	0.05	-0.9119	-0.2330	-1.16E-6
vU0bpstress	1.3881	0.5349	104	2.60	0.0108	0.05	0.3275	2.4488	2.933E-7
varEint	-0.01100	0.2766	104	-0.04	0.9684	0.05	-0.5595	0.5375	-9.49E-7
varEbp mood	0.7228	0.3287	104	2.20	0.0301	0.05	0.07095	1.3746	8.15E-7
varEbp stress	0.6775	0.2250	104	3.01	0.0033	0.05	0.2312	1.1238	2.5E-7

Label	Additional Estimates							
	Estimate	Error	DF	t Value	Pr > t	Alpha	Lower	Upper
Intercept: Men	2.8017	0.3680	104	7.61	<.0001	0.05	2.0719	3.5314
Intercept: Women	2.4604	0.3416	104	7.20	<.0001	0.05	1.7829	3.1379
Intercept: Sex Diff	-0.3413	0.2059	104	-1.66	0.1004	0.05	-0.7495	0.06699
Age Slope: Men	0.06876	0.03007	104	2.29	0.0243	0.05	0.009126	0.1284
Age Slope: Women	-0.01961	0.01506	104	-1.30	0.1955	0.05	-0.04947	0.01024
Age Slope: Sex Diff	-0.08837	0.03327	104	-2.66	0.0092	0.05	-0.1543	-0.02239
BP Mood: Linear	1.6584	0.3974	104	4.17	<.0001	0.05	0.8703	2.4464
Contextual Mood: Linear	1.5560	0.4239	104	3.67	0.0004	0.05	0.7153	2.3967
WP Mood	0.1024	0.1473	104	0.69	0.4887	0.05	-0.1898	0.3945
BP Stress: Men	2.2539	0.5774	104	3.90	0.0002	0.05	1.1089	3.3988
BP Stress: Women	0.7086	0.3137	104	2.26	0.0260	0.05	0.08656	1.3306
BP Stress: Sex Diff	-1.5453	0.6402	104	-2.41	0.0175	0.05	-2.8149	-0.2757
Contextual Stress: Men	2.0496	0.6011	104	3.41	0.0009	0.05	0.8576	3.2415
Contextual Stress: Women	0.6797	0.3329	104	2.04	0.0437	0.05	0.01949	1.3399
Contextual Stress: Sex Diff	-1.3699	0.6707	104	-2.04	0.0436	0.05	-2.6998	-0.03991
WP Stress: Men	0.2043	0.1647	104	1.24	0.2176	0.05	-0.1223	0.5309
WP Stress: Women	0.02886	0.1122	104	0.26	0.7975	0.05	-0.1936	0.2513
WP Stress: Sex Diff	-0.1754	0.1980	104	-0.89	0.3776	0.05	-0.5680	0.2171
Contextual Stress per Day: Men	0.4099	0.1202	104	3.41	0.0009	0.05	0.1715	0.6483
Contextual Stress per Day: Women	0.1359	0.06659	104	2.04	0.0437	0.05	0.003897	0.2680

Pearson Correlation Coefficients, N = 509
Prob > |r| under H0: Rho=0

	symptom5	Pred
symptom5	1.00000	0.52894
symptom5: #Physical Symptoms Truncated at 5		<.0001
Pred	0.52894	1.00000
Predicted Value		<.0001

R = .5289, so R² for the overall model = .28.

It would be impossible to calculate valid pseudo-R² values for each variance component given their heterogeneity across persons and occasions.




Figure 8.3 illustrates the two interactions of stressor days with sex.

Neither sex has a significant WP stress effect (as shown by the slope of the lines), and this non-effect is the same across men and women.

Both men and women show significant effects of reporting more stressor days, even after controlling for current daily stressors. However, these BP and contextual effects of stress are significantly more positive for men (greater difference between lines).